

Physical Examination/Health Record
(to be completed by Family Physician)

All Pupils should have a medical examination before entering Kindergarten. These examinations should include the items on this form. Please take this from to your family physician at the time your child is to be examined. The physician will record findings of physical items, vaccinations, and immunizations required by State Regulations for School enterers.

To be completed by Parent or Health Care Provider	Child's Name: _____	School: TVS: <input type="checkbox"/> ES <input type="checkbox"/> MS <input type="checkbox"/> HS
	Parent's Name: _____	Birth Date: _____ Sex: <input type="checkbox"/> F <input type="checkbox"/> M
	Address: _____	Phone: _____
	_____	HGT: _____ WGT: _____ BP: _____

To be completed by Health Care Provider during and after Physical exam/assessment.	Code	Normal for Age	Abnormal	Not Evaluated
	Eyes			
	Ears			
	Nervous System			
	Teeth			
	Tonsils			
	Adenoids			
	Heart			
	Posture			
	Nutrition			
	Skin			
	Cleanliness			
	Thyroid			
	Genitalia			
Lungs				

Disease History	Yes	No	Date
Measles Rubeola			
Measles Rubella			
Whooping Cough			
Scarlet Fever			
Rheumatic Fever			
Chicken Pox			
Mumps			
Pneumonia			
Polio			
Infectious Hepatitis			
Accident Prone			
Allergies			
Tuberculosis Contact			

Comments: _____

	Vaccine	Dose 1, Date	Dose 2, Date	Dose 3, Date	Dose 4, Date	Dose 5, Date
Immunizations	DtaP, DTP, DT					
	Td					
	Hepatitis B					
	Polio: IPV, OPV	Type:	Type:	Type:	Type:	
	MMR	MMR #1	MMR#2	Measles Only	Mumps Only	Rubella Only
	Hib: Hboc, PRP-OMP, PRT-T PRP-D, COMVAX	Type:	Type:	Type:	Type:	
	Varicella (Chicken Pox)					
Other						

 Physician Signature

 Examination Date

VACCINES	ODH FALL 2019 IMMUNIZATIONS FOR SCHOOL ATTENDANCE
DTaP/DT Tdap/Td Diphtheria, Tetanus, Pertussis	<u>Grade(s) - Kindergarten</u> Four (4) or more of DTaP or DT, or any combination. If all four doses were given before the 4 th birthday, a fifth (5) dose is required. If the fourth dose was administered at least six months after the third dose, and on or after the 4 th birthday, a fifth (5) dose is not required. * <u>Grade(s) - 1-12</u> Four (4) or more of DTaP or DT, or any combination. Three doses of Td or a combination of Td and Tdap is the minimum acceptable for children age seven (7) and up. <u>Grade(s) - 7-12</u> One (1) dose of Tdap vaccine must be administered prior to entry. **
POLIO	<u>Grade(s) - K-9</u> Three (3) or more doses of IPV. The FINAL dose must be administered on or after the 4 th birthday regardless of the number of previous doses. If a combination of OPV and IPV was received, four (4) doses of either vaccine are required. *** <u>Grade(s) 10-12</u> Three (3) or more doses of IPV or OPV. If the third dose of either series was received prior to the fourth birthday, a fourth (4) dose is required; If a combination of OPV and IPV was received, four (4) doses of either vaccine are required.
MMR Measles, Mumps, Rubella	<u>Grade(s) - K-12</u> Two (2) doses of MMR. Dose 1 must be administered on or after the first birthday. The second dose must be administered at least 28 days after dose 1
HEP B Hepatitis B	<u>Grade(s) - K-12</u> Three (3) doses of Hepatitis B. The second dose must be administered at least 28 days after the first dose. The third dose must be given at least 16 weeks after the first dose and at least 8 weeks after the second dose. The last dose in the series (third or fourth dose), must not be administered before age 24 weeks.
Varicella (Chickenpox)	<u>Grade(s) - K-9</u> Two (2) doses of varicella vaccine must be administered prior to entry. Dose 1 must be administered on or after the first birthday. The second dose should be administered at least three (3) months after dose one (1); however, if the second dose is administered at least 28 days after first dose, it is considered valid. <u>Grade(s) - 10-12</u> One (1) dose of varicella vaccine must be administered on or after the first birthday.
MCV4 Meningococcal	<u>Grade(s) 7-10</u> One (1) dose of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry. <u>Grade(s) 12</u> Two (2) doses of meningococcal (serogroup A, C, W, and Y) vaccine must be administered prior to entry****

NOTES:

- Vaccine should be administered according to the most recent version of the Recommended *Immunization Schedules for Children and Adolescents Aged 18 Years or Younger* or the *Catch-up Immunization Schedule for Persons Aged 4 Months Through 18 Years Who Start Late or Who Are More Than 1 Month Behind*, as published by the Advisory Committee on Immunization Practices. Schedules are available for print or download at <http://www.cdc.gov/vaccines/schedules/index.html>
 - Vaccine doses administered ≤ 4 days before the minimum interval or age are valid (grace period). Doses administered ≥ 5 days earlier than the minimum interval or age are not valid doses and should be repeated as age-appropriate. If MMR and Varicella are not given on the same day, the doses must be separated by at least 28 days with no grace period.
 - For additional information please refer to the Ohio Revised Code 3313.67 and 3313.671 for School Attendance and the ODH Director's Journal Entry (available at <https://odh.ohio.gov/wps/portal/gov/odh/know-out-programs/immunizationn/required-vaccines-child-care-school/>) These documents list required and recommended immunizations and indicate exemptions to immunizations.
 - Please contact the Ohio Department of Health Immunization Program at (800) 282-0546 or (614) 466-4643 with questions or concerns.
- * Recommended DTaP or DT minimum intervals for kindergarten students four (4) weeks between doses 1-2 and 2-3; six (6) month minimum intervals between doses 3-4 and 4-5. If a fifth dose is administered prior to the 4th birthday, a sixth dose is recommended but not required.
- ** Pupils who received one dose of Tdap as part of the initial series are not required to receive another dose. Tdap can be given regardless of the interval since the last Tetanus or diphtheria- toxoid containing vaccine. DTaP given to patients age 7 or older can be counted as valid for the one-time Tdap dose.
- *** The final polio dose in the IPV series must be administered at age 4 or older with at least six months between the final and previous dose.
- **** Recommended MCV4 minimum interval of at least eight (8) weeks between dose one (1) and dose two (2). If the first (1st) dose of MCV4 was administered on or after the 16th birthday, a second (2nd) dose is not required. If a pupil is in 12th grade and is 15 years of age or younger, only 1 dose is required. Currently there are no school entry requirements for meningococcal B vaccine.